arXiv:1609.05312v1 [math.AG] 17 Sep 2016

MORDELL-WEIL LATTICE OF INOSE'S ELLIPTIC $K3$ SURFACE ARISING FROM THE PRODUCT OF 3-ISOGENOUS ELLIPTIC CURVES

MASATO KUWATA AND KAZUKI UTSUMI

Abstract. From the product of two elliptic curves, Shioda and Inose [6] constructed an elliptic $K3$ surface having two II^* fibers. Its Mordell-Weil lattice structure depends on the morphisms between the two elliptic curves. In this paper, we give a method of writing down generators of the Mordell-Weil lattice of such elliptic surfaces when two elliptic curves are 3-isogenous. In particular, we obtain a basis of the Mordell-Weil lattice for the singular $K3$ surfaces $X_{[3,3,3]}$, $X_{[3,2,3]}$ and $X_{[3,0,3]}$.

1. Introduction

In the study of the geometry, arithmetic and moduli of $K3$ surfaces, elliptic $K3$ surfaces with large Picard number play a vital role. In 1977 Shioda and Inose [6] gave a classification of singular $K3$ surfaces, that is, $K3$ surfaces with maxium Picard number. For this purpose, they constructed elliptic $K3$ surfaces $\mathcal{E}$ with two singular fibers of type II^* starting from the Kummer surface $Km(E_1 \times E_2)$ with the product of two elliptic curves E_1 and E_2. They constructed $\mathcal{E}$ as a double cover of $Km(E_1 \times E_2)$ with certain properties (now called a Shioda-Inose structure). Later, Inose [1] gave an explicit model of such an elliptic $K3$ surface as a quartic surface in $\mathbf{P}^3$, and remarked that it is the quotient of $Km(E_1 \times E_2)$ by an involution. We call the Kodaira-Néron model of $\mathcal{E}$ the Inose surface associated with E_1 and E_2, and denote it by $Ino(E_1, E_2)$. We thus have a "Kummer sandwich" diagram:

$$Km(E_1 \times E_2) \overset{\pi_2}{\dashrightarrow} Ino(E_1, E_2) \overset{\pi_1}{\dashrightarrow} Km(E_1, E_2)$$

(cf. [7]). Also, $\mathcal{E}$ as an elliptic surface with two II^* fibers is denoted by $F^{(1)}_{E_1,E_2}$. This notation reflects that it is a part of the construction of elliptic $K3$ surfaces of high rank by the first named author [4], where he constructed $F^{(n)}_{E_1,E_2}$, $n = 1, \dots, 6$, which has various Mordell-Weil rank up to 18.

The structure of the Mordell-Weil lattice of $F^{(1)}_{E_1,E_2}$ is known to be isomorphic to $\mathrm{Hom}(E_1, E_2)\langle 2\rangle$ if E_1 and E_2 are nonisomorphic (see [8]). However, given an isogeny $\varphi \in \mathrm{Hom}(E_1, E_2)$ and the Weierstrass equation of $F^{(1)}_{E_1,E_2}$, it is quite difficult to write down the coordinates of the section corresponding to φ, and it has been worked out only in limited cases (cf. [9][2]). Most known examples fall into the case where the degree of isogeny φ equals 2, in which case the calculations are straight forward. One particular example of the case $\deg \varphi = 4$ is dealt in [2, Example 9.2]. In this paper we consider a family of the pairs of elliptic curves E_1 and E_2 with an isogeny $\varphi : E_1 \to E_2$ of degree 3 defined over k. We write down a formula of the section of $F^{(1)}_{E_1,E_2}$ coming from φ defined over the base field k. To do so, we first work with the surface $F^{(6)}_{E_1,E_2}$, which has a simple affine model that can be viewed as a family of cubic curves with a rational point over k. We modify the method

2010 *Mathematics Subject Classification.* Primary 14J27, 14J28; Secondary 14H52, 11G05.
Key words and phrases. $K3$ surface, elliptic surface, elliptic curve.

in [2] to find sections of $F^{(1)}_{E_1,E_2}$. We also give a section of $F^{(2)}_{E_1,E_2}$ coming from the isogeny φ, and give a basis defined over the field $k(E_1[2], E_2[2])$ when E_1 and E_2 do not have a complex multiplication.

In §7 we study some examples of singular $K3$ surfaces in detail. In particular, we determine a basis of the MWL of the Inose surface $F^{(1)}_{E_1,E_2}$ and that of $F^{(2)}_{E_1,E_2}$ for the singular $K3$ surfaces $X_{[3,3,3]}$, $X_{[3,2,3]}$ and $X_{[3,0,3]}$ which correspond to the quadratic form $3x^2+3xy+3y^2$, $3x^2+2xy+3y^2$, and $3x^2+3y^2$ respectively.

Acknowledgements. We would like to express sincere gratitude to Professor Ichiro Shimada for useful discussions. We would also like to thank Professor Hisanori Ohashi for his helpful comments. The computer algebra system `Maple` was used in the calculations for this paper. Kuwata was partially supported by JSPS Grant-in-Aid for Scientific Research (C) No. 26400023, and by the Chuo University Grant for Special Research.

2. Inose surface

Throughout this paper the base field k of algebraic varieties is assumed to be a number field.

Let $Km(E_1 \times E_2)$ be the Kummer surface associated with the product of elliptic curves E_1 and E_2, that is, the minimal resolution of the quotient surface $E_1 \times E_2/\{\pm 1\}$. If E_1 and E_2 are defined by the equations

$$E_1 : y_1^2 = f_1(x_1), \quad E_2 : y_2^2 = f_2(x_2),$$

where $f_1(x_1)$ and $f_2(x_2)$ are cubic polynomials, an affine singular model of $Km(E_1 \times E_2)$ is given as the hypersurface in $\mathbf{A}^3$ defined by the equation

$$f_2(x_2) = t^2 f_1(x_1), \tag{2.1}$$

where $t = y_2/y_1$. Then the map $Km(E_1 \times E_2) \to \mathbf{P}^1$ induced by $(x_1, x_2, t) \mapsto t$ is a Jacobian fibration, which is sometimes called Inose's pencil (cf. [5]).

Take a parameter u such that $t = u^3$, and consider (2.1) as a family of cubic curves in $\mathbf{P}^2$, or a cubic curve over $k(u)$. Then, we see that it has a rational point $(1 : u^2 : 0)$. Using this point as the origin, we consider it as an elliptic curve over $k(u)$. In particular, if $f_1(x_1)$ and $f_2(x_2)$ are given by

$$f_1(x_1) = x_1^3 + a_2x_1^2 + a_4x_1 + a_6, \quad \text{and} \quad f_2(x_2) = x_2^3 + a_2'x_2^2 + a_4'x_2 + a_6',$$

then we can convert (2.1) to the Weierstrass form

$$Y^2 = X^3 - \frac{1}{3}A\,X + \frac{1}{64}\Big(\Delta_{E_1}u^6 + B + \frac{\Delta_{E_2}}{u^6}\Big),$$

where

$$\begin{cases} A = (a_2^2 - 3a_4)({a_2'}^2 - 3a_4'), \\ B = \dfrac{32}{27}(2a_2^3 - 9a_2a_4 + 27a_6)(2{a_2'}^3 - 9a_2'a_4' + 27a_6'), \\ \Delta_{E_1} = 16(a_2^2a_4^2 - 4a_2^3a_6 + 18a_2a_4a_6 - 4a_4^3 - 27a_6^2), \\ \Delta_{E_2} = 16({a_2'}^2{a_4'}^2 - 4{a_2'}^3a_6' + 18a_2'a_4'a_6' - 4{a_4'}^3 - 27{a_6'}^2). \end{cases}$$

Let $s = t^2 = u^6$. Define $F^{(1)}_{E_1E_2}$ to be

$$F^{(1)}_{E_1E_2} : Y^2 = X^3 - \frac{1}{3}A\,X + \frac{1}{64}\Big(\Delta_{E_1}s + B + \frac{\Delta_{E_2}}{s}\Big). \tag{2.2}$$

This Jacobian fibration has two reducible fibers of type II^* at $s = 0$ and $s = \infty$.

Definition 2.1 (cf. [2])**.** The Kodaira-Néron model of the elliptic surface $F^{(1)}_{E_1,E_2}$ over k defined by (2.2) is called the Inose surface associated with E_1 and E_2, and it is denoted by $Ino(E_1, E_2)$.

Remark 2.2. Over a certain extension of k, the equation of $F^{(1)}_{E_1,E_2}$ may be given by

$$F^{(1)}_{E_1E_2} : Y^2 = X^3 - 3\sqrt[3]{J_1J_2}\, X + s + \frac{1}{s} - 2\sqrt{(1-J_1)(1-J_2)},$$

where $J_i = j(E_i)/1728$ $(i = 1, 2)$ is the absolute j-invariant of E_i (cf. [1][8]).

Definition 2.3. For $n \geq 1$, the elliptic surface $F^{(n)}_{E_1,E_2}$ over k is defined by

$$F^{(n)}_{E_1E_2} : Y^2 = X^3 - \frac{1}{3}A\, X + \frac{1}{64}\Big(\Delta_{E_1}s^n + B + \frac{\Delta_{E_2}}{s^n}\Big). \tag{2.3}$$

Remark 2.4. (1) The Kodaira-Néron model of $F^{(n)}_{E_1,E_2}$ is a $K3$ surface for $n = 1, \ldots, 6$, but not for $n \geq 7$ ([4]).

(2) Since $u^6 = t^2$, Inose's pencil on $Km(E_1 \times E_2)$ is isomorphic to $F^{(2)}_{E_1,E_2}$. However, the isomorphism between (2.1) and (2.3) for $n = 2$ may not be defined over k itself.

3. Mordell-Weil lattice of the Inose surface

In this section we give a summary of known facts on the Mordell-Weil lattice of $F^{(1)}_{E_1,E_2}$ over $\bar{k}(s)$.

Theorem 3.1. [8, Theorem 6.3] *The Jacobian fibration $F^{(1)}_{E_1,E_2}$ on $Ino(E_1, E_2)$ has two singular fibers of type* II^* *at $s = 0$ and ∞, and the other singular fibers and its Mordell-Weil rank are given in the table below. Here, h is the rank of* $\mathrm{Hom}_{\bar{k}}(E_1, E_2)$.

j-invariants	*singular fibers*	*Mordell-Weil rank*
$J_1 \neq J_2, J_1J_2 \neq 0$	$4\,\mathrm{I}_1$	h
$J_1 \neq J_2, J_1J_2 = 0$	$2\,\mathrm{II}$	h
$J_1 = J_2 \neq 0, 1$	$\mathrm{I}_2, 2\,\mathrm{I}_1$	$h-1$
$J_1 = J_2 = 1$	$2\,\mathrm{I}_2$	0
$J_1 = J_2 = 0$	IV	0

Assume $J_1 \neq J_2$. Then the Mordell-Weil group $F^{(1)}_{E_1,E_2}(\overline{\mathbf{Q}}(s))$ is torsion-free, and isomorphic to the lattice $\mathrm{Hom}_{\bar{k}}(E_1, E_2)\langle 2\rangle$, *where the pairing of* $\mathrm{Hom}_{\bar{k}}(E_1, E_2)$ *is given by*

$$(\varphi, \psi) = \frac{1}{2}\left(\deg(\varphi + \psi) - \deg\varphi - \deg\psi\right) \quad \varphi, \psi \in \mathrm{Hom}_{\bar{k}}(E_1, E_2).$$

Remark 3.2. The notation $\langle n\rangle$ means that the paring of the lattice is multiplied by n.

4. 3-isogenies of elliptic curves

We recall some general facts on 3-isogenies between elliptic curves following J. Top [10, §3].

Let E be an elliptic curve over a number field k, and $G \subset E(\bar{k})$ a subgroup of order three that is stable under the action of $\mathrm{Gal}(\bar{k}/k)$. E can be given by an equation of the form $y^2 = f(x)$ with $\deg f = 3$. Then G consists of three points $O = (0 : 1 : 0)$, $P = (\alpha, \beta)$, and $2P = -P = (\alpha, -\beta)$. Since G is Galois invariant, we have $\alpha \in k$ and $\beta^2 \in k$. Replacing x by $x + \alpha$ if necessary, we may assume $\alpha = 0$. It turns out that we have two cases. Either (i) E is given by $y^2 = x^3 + d$ and $P = (0, \sqrt{d})$, or (ii) E is given by $y^2 = x^3 + a(x - b)^2$ and $P = (0, b\sqrt{a})$.

If E is given by $y_1^2 = x_1^3 + d$ and $P = (0, \sqrt{d})$, then the quotient E/G is given by the equation

$$E/G : y_2^2 = x_2^3 - 27d,$$

and the quotient map $\varphi : E \to E/G$ is given by

$$\varphi : (x_1, y_1) \mapsto (x_2, y_2) = \left(\frac{x_1^3 + 4d}{x_1^2}, \frac{(x_1^3 - 8d)y_1}{x_1^3} \right).$$

In this case E and E/G are isomorphic over $\bar{k}$, and the j-invariants are equal to 0.

If E is given by

$$E : y_1^2 = x_1^3 + a(x_1 - b)^2,$$

and $P = (0, b\sqrt{a})$, then the quotient E/G is given by

$$\hat{E} = E/G : y_2^2 = x_2^3 - 27a(x_2 - 4a - 27b)^2.$$

The isogeny $\varphi : E \to E/G$ is given by $\varphi_1(x_1, y_1) = (\varphi_x(x_1), \varphi_y(x_1)y_1)$, where

$$\varphi_x(x_1) = \frac{3\left(3x_1^3 + 4ax_1^2 - 12abx_1 + 12ab^2\right)}{x_1^2}, \quad \varphi_y(x_1) = -\frac{27\left(x_1^3 + 4abx_1 - 8ab^2\right)}{x_1^3}.$$

5. The rational section of $F^{(1)}_{E_1,E_2}$ arising from a 3-isogeny

In this section, we assume E_1 and E_2 are 3-isogenous over k, and we find an explicit section of $F^{(1)}_{E_1,E_2}$.

In the case where the j-invariants of both E_1 and E_2 are equal to 0, the Mordell-Weil group $F^{(1)}_{E_1,E_2}(\bar{k}(s))$ is trivial by Theorem 3.1, and we have nothing to do.

As for the second case in §4, we modify the equations slightly by replacing x_2 by $3x_2$, and y_2 by $27y_2$, so that E_1 and E_2 are given by

$$\begin{aligned} E_1 &: y_1^2 = x_1^3 + a(x_1 - b)^2, \\ E_2 &: 27y_2^2 = x_2^3 - a(3x_2 - 4a - 27b)^2, \end{aligned} \tag{5.1}$$

where $a, b \in k$. We work with the cubic curve over $k(u)$ in $\mathbf{P}^2 = \{(x_1 : x_2 : z)\}$ given by

$$C_u : x_2^3 - a(3x_2 - 4az - 27bz)^2 z = 27u^6\big(x_1^3 + a(x_1 - bz)^2 z\big), \tag{5.2}$$

which is isomorphic over k to $F^{(6)}_{E_1,E_2}$ with the choice of origin $O = (1 : 3u^2 : 0)$. Its Weierstrass equation is given by

$$\begin{aligned} F^{(6)}_{E_1,E_2} : Y^2 = X^3 &- \frac{16}{3}a^2\big(a^2 - 12bb'\big)X \\ &- 16a^2bb'\Big(b^2u^6 + \frac{b'^2}{u^6}\Big) + \frac{32}{27}a^2\big(4a^4 + 144a^2bb' - 27b^2b'^2\big), \end{aligned} \tag{5.3}$$

where $b' = 4a + 27b$.

Remark 5.1. The origin O in our case is not an inflection point of the cubic curve C_u. Thus, three collinear points P, Q, R do not satisfy the equation $P+Q+R = O$ under the group law. Instead, we have $P + Q + R = \overline{O}$, where $\overline{O}$ is the third point of intersection between C_u and the tangent line at O.

Remark 5.2. By definition, the surface $F^{(1)}_{E_1,E_2}$ is obtained as the quotient of (5.3) by the action $(X, Y, u) \mapsto (X, Y, -\omega u)$, where ω is a third root of unity. However, the action $\big((x_1 : x_2 : z), u\big) \mapsto \big((x_1 : x_2 : z), -\omega u\big)$ on C_u does not correspond to this action since the quotient of the latter gives a rational surface. As a matter of fact, it turns out that the action $\big((x_1 : x_2 : z), u\big) \mapsto \big((x_1 : x_2 : z), -\omega u\big)$ corresponds to the action $(X, Y, u) \mapsto (X, -Y, -\omega u)$.

With our choice of coordinates, we have $\varphi_x(x_1) = (3x_1^3 + 4ax_1^2 z - 12abx_1 z^2 + 12ab^2 z^3)/x_1^2 z$. Replacing x_2 by $\varphi_x(x_1)$ in (5.2), we have

$$\frac{27}{x_1^6}\big(x_1^3 + a(x_1 z - bz)^2 z\big)\left((x_1^3 + 4abx_1 z^2 - 8ab^2 z^3)^2 - (u^3 x_1^3)^2\right) = 0.$$



FIGURE 1. The x_1-coordinate of $Q_i^{\pm}$ $(i = 1, 2, 3)$ is a solution of the cubic equation $\varphi_y(x_1) = \pm u^3$ in $\overline{k(u)}$. A conic $\mathcal{Q}^+$, which passes Q_1^+, Q_2^+, Q_3^+, and tangent to to the cubic C_u (5.2) at O, intersects with C_u at the sixth point Q_4^+.

Define two homogeneous polynomials in x_1, x_2, z

$$p^+(x_1, z) = x_1^3 + 4abx_1z^2 - 8ab^2z^3 - u^3x_1^3,$$
$$p^-(x_1, z) = x_1^3 + 4abx_1z^2 - 8ab^2z^3 + u^3x_1^3,$$

and let $D_\varphi^+ = Q_1^+ + Q_2^+ + Q_3^+$ be the divisor on C_u of degree 3 such that the x_1-coordinates of Q_1^+, Q_2^+, Q_3^+ are the three roots of $p^+(x_1, z) = 0$. We look for a conic $\mathcal{Q}$ given by

$$\mathcal{Q} : q(x_1, x_2, z) = a_1x_1^2 + a_2x_1x_2 + a_3x_2^2 + a_4x_1z + a_5x_2z + a_6z^2 = 0 \tag{5.4}$$

which passes through Q_1^+, Q_2^+, Q_3^+, and tangent to the tangent line $\mathcal{T}$ of C_u at the origin O (see Figure 1).

The tangent line $\mathcal{T}$ is given by

$$\mathcal{T} : l(x_1, x_2, z) = 3u^2x_1 - x_2 + a(u^2 + 3)z = 0. \tag{5.5}$$

The condition that $\mathcal{T}$ is tangent to $\mathcal{Q}$ is obtained as follows. Regard $q(x_1, x_2, z)$ and $l(x_1, x_2, z)$ as polynomials in x_2, and compute the remainder of $q(x_1, x_2, z)$ divided by $l(x_1, x_2, z)$. Then, $\mathcal{T}$ is tangent to $\mathcal{Q}$ at O if and only if the coefficients of x_1z and x_1^2 vanish. Similarly, $\mathcal{Q}$ passes through Q_1^+, Q_2^+, Q_3^+ if and only if $q(x_1, \varphi_x(x_1), z)$ is divisible by $p^+(x_1, z)$. We then obtain a system of five homogeneous linear equations in $a_1, \ldots, a_6$. It is easy to solve this system to find a conic $\mathcal{Q}$. By Bźout's theorem, the intersection $\mathcal{Q}$ and C_u consists of six points counting multiplicity. It is not difficult to find the sixth point Q_4^+, but the result is rather complicated to show here.

Similarly, we obtain Q_4^- starting from the divisor $D_\varphi^- = Q_1^- + Q_2^- + Q_3^-$ obtained from $p^-(x_1, z)$.

Proposition 5.3. *Let D_φ^+ (resp. D_φ^-) be the divisor on the cubic curve C_u defined by the equation $\varphi_y(x_1) = u^3$ (resp. $\varphi_y(x_1) = -u^3$).*

(i) *The divisor D_φ^+ (resp. D_φ^-) determines a k-rational point P_φ^+ (resp. P_φ^-) in $F^{(6)}_{E_1,E_2}(k(u))$.*

(ii) *$P_\varphi^+ - P_\varphi^-$ is in the image of $F^{(1)}_{E_1,E_2}(k(s)) \to F^{(6)}_{E_1,E_2}(k(u))$ induced by $s \mapsto u^6$. The height of its pre-image in $F^{(1)}_{E_1,E_2}(k(s))$ is 6.*

Proof. Since D_φ^+ and D_φ^- are both defined over $k(u)$, Q_4^+ and Q_4^- are $k(u)$-rational points on C_u. We identify $C_u(\bar{k}(u))$ with $\mathrm{Pic}^0_{\bar{k}}(C_u) \simeq F^{(6)}_{E_1,E_2}(\bar{k}(u))$ via the map $P \mapsto [P - O]$, where $[\ \ ]$ stands for the divisor class. Let P_φ^+ (resp. P_φ^-) be the point in $F^{(6)}_{E_1,E_2}(k(u))$ corresponding to Q_4^+ (resp. Q_4^-) in $C_u(k(u))$, or equivalently, $P_\varphi^\pm = [Q_4^\pm - O]$. Let σ be the automorphism $\big((x_1 : x_2 : z), u\big) \mapsto \big((x_1 : x_2 : z), -\omega u\big)$ on C_u. As we noted in Remark 5.2, the corresponding automorphism $\tilde{\sigma}$ on $F^{(6)}_{E_1,E_2}$ sends (X, Y, u) to $(X, -Y, -\omega u)$. Clearly, the involution σ^3 interchanges between Q_4^+ and Q_4^-, and thus $([-1] \circ \tilde{\sigma})^3 : (X, Y, u) \mapsto (X, -Y, -u)$ leaves $P_\varphi^+ - P_\varphi^-$ invariant. Moreover, since $Q_4^\pm$ are both invariant under the automorphism σ^2 by construction, $P_\varphi^+ - P_\varphi^-$ is also invariant under $([-1]\circ\tilde{\sigma})^2$. This implies that $P_\varphi^+ - P_\varphi^-$ is invariant under $[-1] \circ \tilde{\sigma}$ and it belongs to the image of $F^{(1)}_{E_1,E_2}(k(s))$ under the map $s \mapsto u^6$. The calculation of the height is the same as in [8]. □

We denote by $P_\varphi^{(1)}$ the section $P_\varphi^+ - P_\varphi^-$ in $F^{(1)}_{E_1,E_2}(k(s))$. Carrying out calculations according to the above recipe, we obtain the coordinates of $P_\varphi^{(1)}$ as follows:

$$P_\varphi^{(1)} = \left(\frac{\big(3S^3 + c_2S^2 + c_1S + c_0\big)}{48a^2\big(S - 2bb'\big)}, \frac{-s\big(bs + b'\big)\big(S^4 + d_3S^3 + d_2S^2 + d_1S + d_0\big)}{64a^3\big(bs - b'\big)^3} \right),$$

where

$$\begin{aligned}
S &= b^2 s + \frac{b'^2}{s}, \\
c_2 &= -6\big(8a^2 - 3bb'\big), \quad c_1 = 4\big(16a^4 + 48a^2bb' + 9b^2b'^2\big), \\
c_0 &= 8bb'\big(80a^4 + 72a^2bb' + 3b^2b'^2\big), \\
d_3 &= -8\big(3a^2 - bb'\big), \quad d_2 = 8(16a^4 + 6a^2bb' + 3b^2b'^2), \\
d_1 &= -32bb'\big(24a^4 - 15a^2bb' - b^2b'^2\big), \\
d_0 &= 16bb'\big(128a^6 + 256a^4bb' + 36a^2b^2b'^2 + b^3b'^3\big).
\end{aligned}$$

6. Rational sections of $F^{(2)}_{E_1,E_2}$

Next, we consider the Mordell-Weil lattice $F^{(2)}_{E_1,E_2}$. We use the same notation as the previous section.

Let $\alpha_1, \alpha_2, \alpha_3$ be the three roots of $x_1^3 + a(x_1 - b)^2 = 0$. For $i = 1, 2, 3$, define $\beta_i = \varphi_x(\alpha_i)$. Since φ is an isogeny of odd degree, $(\beta_i, 0)$ are the 2-torsion points of E_2. Thus, we have $k(E_1[2], E_2[2]) = k(\alpha_1, \alpha_2, \alpha_3)$, which we denote by k_2 for short.

Denote by R_{ij} be the point $(\alpha_i : \beta_j : 1)$ in C_u. The quotient of C_u by the action $u \mapsto \omega u$ is nothing but the model of Kummer surface (2.1). It is known (cf. [2, Prop. 3.3]) that, by taking R_{11} as the origin, $F^{(2)}_{E_1,E_2}(\bar{k}(t))$ is generated by $R_{22}, R_{33}, R_{23}, R_{32}$.

Let O_ω and O_{ω^2} be the points in C_u defined by $\big(1 : 3(\omega u)^2 : 0\big)$ and $\big(1 : 3(\omega^2 u)^2 : 0\big)$ respectively. Clearly, the action $u \mapsto \omega u$ induces a cyclic permutation $O \mapsto O_\omega \mapsto O_{\omega^2}$. Thus, the divisor $O + O_\omega + O_{\omega^2}$ is invariant under this action. Let $q(x_1, x_2, z)$ be the quadratic form in (5.4) and $l(x_1, x_2, z)$ the linear form in (5.5). Consider the function $f = \frac{q(x_1,x_2,z)}{z\, l(x_1,x_2,z)}$. The divisor of this function is

$$\mathrm{div}(f) = D_\varphi^+ + Q_4^+ + 2O - (O + O_\omega + O_{\omega^2} + 2O + \overline{O}).$$

Thus, we have

$$Q_4^+ - \overline{O} \sim (O + O_\omega + O_{\omega^2}) - D_\varphi^+.$$

Let $P_{\overline{O}}$ be the section in $F^{(6)}_{E_1,E_2}(k(u))$ corresponding to $\overline{O} \in C_u(k(u))$. Then, since $(O + O_\omega + O_{\omega^2}) - D_\varphi^+$ is invariant under the action $u \mapsto \omega u$, the section $P_\varphi^+ - P_{\overline{O}}$ is also invariant under this action. This implies that $P_\varphi^+ - P_{\overline{O}}$ is a $k(t)$-rational section where $t = u^3$.

Proposition 6.1. *Let $P_\varphi^{(2)} = P_\varphi^+ - P_{\overline{O}}$. Then $P_\varphi^{(2)}$, R_{22}, R_{33}, R_{23}, R_{32} are linearly independent. They generate the Mordell-Weil group $F^{(2)}_{E_1,E_2}(\bar{k}(t))$ if E_1 and E_2 do not have complex multiplication.*

Proof. By straightforward calculations, we have

$$P_\varphi^{(2)} = \left(T^2 + 4aT + \frac{4}{3}(a^2 - 3bb'), \left(bt + \frac{b'}{t}\right)\big(T^2 + 6aT + 4(2a^2 + bb')\big) \right),$$

where $b' = 4a + 27b$ and $T = bt - b'/t$. Since the coordinates of R_{ij} involve the roots of the cubic equation $x_1^3 + a(x_1 - b)^2 = 0$, we do not write down the explicit coordinates here, but calculations are straightforward. (We will show them in the numerical examples.) The height matrix with respect to $P_\varphi^{(2)}$, R_{22}, R_{33}, R_{23}, R_{32} is given by

$$\frac{1}{3}\begin{pmatrix} 12 & 0 & 0 & -3 & -3 \\ 0 & 4 & 2 & 0 & 0 \\ 0 & 2 & 4 & 0 & 0 \\ -3 & 0 & 0 & 4 & 2 \\ -3 & 0 & 0 & 2 & 4 \end{pmatrix}.$$

Its determinant is $2^4/3$ as required in [8, Theorem 1.2] □

Remark 6.2. The height paring of given two rational points is easily computed by the height formula in [3].

7. Singular $K3$ surfaces

A complex $K3$ surface whose Picard number equals to maximum possible number 20 is called a singular $K3$ surface. Shioda and Inose showed that a complex singular $K3$ surface is isomorphic to an Inose surface $Ino(E_1, E_2)$ for some elliptic curves E_1 and E_2 that have complex multiplication and are isogenous to each other.

Theorem 7.1 ([6])**.** *There is a one-to-one correspondence between the set of isomorphism classes of complex singular $K3$ surfaces and the set of equivalence classes of positive-definite even integral lattices of rank 2 with respect to $SL_2(\mathbf{Z})$:*

$$\begin{aligned} &\{ \text{ singular } K3 \text{ surfaces over } \overline{\mathbf{Q}} \} \,/\, \text{isom.} \\ &\overset{1:1}{\leftrightarrow} \left\{ \begin{pmatrix} 2a & b \\ b & 2c \end{pmatrix} \;\middle|\; a, b, c \in \mathbf{Z},\ a, c > 0,\ b^2 - 4ac < 0 \right\} \Big/ SL_2(\mathbf{Z}), \end{aligned}$$

which associates a singular $K3$ surface X with its transcendental lattice T_X.

In fact, a singular $K3$ surface corresponding to the lattice $Q = \begin{pmatrix} 2a & b \\ b & 2c \end{pmatrix}$ is constructed as follows. Let E_1 and E_2 be the complex elliptic curves $\mathbf{C}/\mathbf{Z} \oplus \tau_1\mathbf{Z}$ and $\mathbf{C}/\mathbf{Z} \oplus \tau_2\mathbf{Z}$, respectively, where

$$\tau_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}, \quad \tau_2 = \frac{b + \sqrt{b^2 - 4ac}}{2}.$$

Then, the Inose surface $Ino(E_1, E_2)$ is a singular $K3$ surface corresponding to Q, which is often denoted by $X_{[a,b,c]}$.

In the following, we study in detail the Mordell-Weil group of $F^{(1)}$ and $F^{(2)}$ for $X_{[3,3,3]}$, $X_{[3,2,3]}$, and $X_{[3,0,3]}$

7.1. **The singular $K3$ surface $X_{[3,3,3]}$.** The transcendental lattice of the singular $K3$ surface $X_{[3,3,3]}$ is given by the matrix

$$\begin{pmatrix} 6 & 3 \\ 3 & 6 \end{pmatrix}.$$

Then, elliptic curves E_1 and E_2 are given by

$$E_1 = \mathbf{C}/\mathbf{Z} \oplus \mathbf{Z}\omega, \quad E_2 = \mathbf{C}/\mathbf{Z} \oplus \mathbf{Z}(-3\omega^2),$$

where $\omega = \frac{-1+\sqrt{-3}}{2}$. The j-invariants are given by

$$j(E_1) = 0, \quad j(E_2) = -1288000,$$

and we take the following Weierstrass forms of them:

$$\begin{aligned} E_1 : \quad y_1^2 &= x_1^3 + 6(x_1 + 1)^2, \\ E_2 : 27y_2^2 &= x_2^3 - 6(3x_2 + 3)^2 \end{aligned}$$

This is nothing but the case $a = 6$, $b = -1$ in (5.1). Then, $F^{(1)}_{E_1,E_2}$ is given by

$$F^{(1)}_{E_1,E_2} : Y^2 = X^3 - 27\Big(s - 506 + \frac{9}{s}\Big),$$

after rescaling X and Y. In this case, $F^{(1)}_{E_1,E_2}$ itself has complex multiplication by ω.

There are two 3-isogenies from E_1 to E_2. One is the 3-isogeny φ in §4. We denote it by φ_1. The other, denoted by φ_2, is a composite of φ_1 and multiplication by ω on E_2. They are given by

$$\varphi_1(x_1, y_1) = (\varphi_x(x_1), \varphi_y(x_1)y_1), \quad \varphi_2(x_1, y_1) = (\omega\varphi_x(x_1), \varphi_y(x_1)y_1),$$

where

$$\varphi_x(x_1) = \frac{3(x_1^3 + 8x_1^2 + 24x_1 + 24)}{x_1^2}, \quad \varphi_y(x_1) = -\frac{x_1^3 - 24x_1 - 48}{x_1^3}.$$

The isogeny φ_1 yields a $\mathbf{Q}$-rational section $P^{(1)}_\varphi = P^+_\varphi - P^-_\varphi$ of $F^{(1)}_{E_1,E_2}$ (see §5). If we let $S = \frac{1}{2}\big(\frac{s}{3} + \frac{3}{s}\big)$, its coordinates are given by

$$P^{(1)}_\varphi = \Bigg(\frac{S^3 - 93S^2 + 963S + 4129}{64(S-1)}, \\ \frac{3s(s+3)(S^4 - 140S^3 + 4758S^2 - 13100S + 258481)}{256(s-3)^3}\Bigg).$$

The sections $P^{(1)}_\varphi$ and $[-\omega]P^{(1)}_\varphi = \big(\omega X, -Y\big)$ form a basis of the Mordell-Weil group $F^{(1)}_{E_1,E_2}(\overline{\mathbf{Q}}(s)) = F^{(1)}_{E_1,E_2}(\mathbf{Q}(\omega)(s))$. The height matrix with respect to this basis is given by $\begin{pmatrix} 6 & 3 \\ 3 & 6 \end{pmatrix}$ as expected.

Next, we consider $F^{(2)}_{E_1,E_2}(\overline{\mathbf{Q}}(t))$. The explicit formula for the sections $P^{(2)}_{\varphi}$, R_{22}, R_{33}, R_{23}, R_{32} described in Proposition 6.1 are as follows:

$$P^{(2)}_{\varphi} = \left(\frac{1}{4}T_-^2 - 6T_- + 15, \frac{1}{8}T_+\big(T_-^2 - 36T_- + 300\big)\right)$$
$$R_{22} = \left(-15\cdot 2^{\frac{2}{3}}\omega, -3\sqrt{-3}\,T_-\right), \quad R_{33} = [-\omega]R_{22},$$
$$R_{23} = \left(-24\omega, -3\sqrt{-3}\,T_+\right), \quad R_{32} = [-\omega]R_{23},$$

where $T_- = t - 3/t$, and $T_+ = t + 3/t$.

The height matrix with respect to $P^{(2)}_{\varphi}$, $[-\omega]P^{(2)}_{\varphi}$,$R_{22}$, R_{33}, R_{23}, R_{32} is given by

$$\frac{1}{3}\begin{pmatrix} 12 & 6 & 0 & 0 & -3 & -3 \\ 6 & 12 & 0 & 0 & 0 & -3 \\ 0 & 0 & 4 & 2 & 0 & 0 \\ 0 & 0 & 2 & 4 & 0 & 0 \\ -3 & 0 & 0 & 0 & 4 & 2 \\ -3 & -3 & 0 & 0 & 2 & 4 \end{pmatrix}.$$

Its determinant is $2^2 \cdot 3$ as expected.

7.2. **The singular $K3$ surface $X_{[3,2,3]}$.** The transcendental lattice of the singular $K3$ surface $X_{[3,2,3]}$ is given by the matrix

$$\begin{pmatrix} 6 & 2 \\ 2 & 6 \end{pmatrix}.$$

Then, elliptic curves E_1 and E_2 are given by $\mathbf{C}/\mathbf{Z} \oplus \tau_1\mathbf{Z}$ and $\mathbf{C}/\mathbf{Z} \oplus \tau_2\mathbf{Z}$, where

$$\tau_1 = \frac{-1 + 2\sqrt{-2}}{3}, \quad \tau_2 = 1 + 2\sqrt{-2}.$$

The j-invariants of E_1 and E_2 are given by

$$j(E_1) = 26125000 - 18473000\sqrt{2}, \quad j(E_2) = 26125000 + 18473000\sqrt{2}.$$

We thus work on the base field $k = \mathbf{Q}(\sqrt{2})$. We choose E_1 and E_2 such that their Weierstrass forms are given by

$$E_1 : y_1^2 = x_1^3 + 6(3 - \sqrt{2})x_1^2 + 9(3 + 2\sqrt{2} + 3)x_1,$$
$$E_2 : y_2^2 = x_2^3 + 6(3 + \sqrt{2})x_2^2 + 9(3 - 2\sqrt{2} + 3)x_2.$$

They are so-called $\mathbf{Q}$-curves; they are Galois conjugates and isogenous to each other.

Both E_1 and E_2 have complex multiplication by $\mathbf{Z}[1+2\sqrt{-2}]$. Let $K = \mathbf{Q}(\sqrt{-2})$, and let $H = kK = \mathbf{Q}(\sqrt{2}, \sqrt{-2})$ be the Hilbert class field of K. They admit two isogenies $\varphi_1, \varphi_2 : E_1 \to E_2$ of degree 3 defined over H given by

$$\varphi_1(x_1, y_1) = (\varphi_{1,x}(x_1), \varphi_{1,y}(x_1)y_1), \quad \varphi_2(x_1, y_1) = (\varphi_{2,x}(x_1), \varphi_{2,y}(x_1)y_1),$$

where

$$\varphi_{1,x}(x_1) = \frac{(1 - 2\sqrt{-2})x_1(x_1 + 3 - 3\sqrt{-2})^2}{9(x_1 + 3 + 3\sqrt{-2} + 2\sqrt{2} + 4\sqrt{-1})^2},$$
$$\varphi_{2,x}(x_1) = \frac{(1 + 2\sqrt{-2})x_1(x_1 + 3 + 3\sqrt{-2})^2}{9(x_1 + 3 - 3\sqrt{-2} + 2\sqrt{2} - 4\sqrt{-1})^2}.$$

The isogeny φ_i yields a H-rational section $P^+_{\varphi_i}$ of $F^{(6)}_{E_1,E_2}$ for $i = 1, 2$.

Let $\hat{\varphi}_i : E_2 \to E_1$ be the dual isogeny of φ_i for $i = 1, 2$. The endomorphisms $\hat{\varphi}_2 \circ \varphi_1$ and $\hat{\varphi}_1 \circ \varphi_2$ correspond to the complex multiplication of E_1 by $1 \pm 2\sqrt{-2}$.

The surface $F^{(6)}_{E_1,E_2}$ is defined over k and given by

$$F^{(6)}_{E_1,E_2} : Y^2 = X^3 + \frac{575}{12}X + \Big(\frac{u^6}{(1-\sqrt{2})^3} - \frac{34937}{108} - \frac{(1-\sqrt{2})^3}{u^6}\Big).$$

By letting $s' = s/(1-\sqrt{2})^3 = u^{1/6}/(1-\sqrt{2})^3$, the Weierstrass equation of the surface $F^{(1)}_{E_1,E_2}$ is given by

$$F^{(1)}_{E_1,E_2} : Y^2 = X^3 + \frac{575}{12}X + \Big(s' - \frac{34937}{108} - \frac{1}{s'}\Big).$$

With this model, the X-coordinate of the section $P^{(1)}_{\varphi_1} = P^+_{\varphi_1} - P^-_{\varphi_1}$ of $F^{(1)}_{E_1,E_2}$ (see §5) is given by

$$X(P^{(1)}_{\varphi_1}) = \frac{-\sqrt{-1}}{12(2-\sqrt{-1})^8(S+2\sqrt{-1})}(3S'^3 + c_2S'^2 + c_1S' + c_0),$$

where

$$S' = s' - \frac{1}{s'}, \quad c_2 = -42(23-10\sqrt{-1}), \quad c_1 = 2(9402-13685\sqrt{-1}),$$
$$c_0 = -4(61663+50160\sqrt{-1}).$$

The Y-coordinate can be obtained easily, but it is rather complicated and we do not include here. The section $P^{(1)}_{\varphi_2}$ is the image of $P^{(1)}_{\varphi_1}$ under the complex conjugate $\sqrt{-1} \mapsto -\sqrt{-1}$. The sections $P^{(1)}_{\varphi_1}$ and $P^{(1)}_{\varphi_2}$ form a basis of the Mordell-Weil group $F^{(1)}_{E_1,E_2}(\bar{k}(s)) = F^{(1)}_{E_1,E_2}(H(s))$. The height matrix with respect to this basis is given by $\begin{pmatrix} 6 & 2 \\ 2 & 6 \end{pmatrix}$ as expected.

Let $L = H\big(\sqrt{1-\sqrt{2}}\big) = \mathbf{Q}\big(\sqrt{1-\sqrt{2}}, \sqrt{-1}\big)$. Then, all the 2-torsion points of E_1 and E_2 are defined over L, and thus, the field of definition of $F^{(2)}_{E_1,E_2}(\bar{k}(t))$ is L. If we let $t' = u^{1/3}/(1-\sqrt{2})^{3/2}$, the Weierstrass equation of $F^{(2)}_{E_1,E_2}$ is given by

$$F^{(2)}_{E_1,E_2} : Y^2 = X^3 + \frac{575}{12}X + \Big(t'^2 - \frac{34937}{108} - \frac{1}{t'^2}\Big).$$

Let $P^{(2)}_{\varphi_i} = P^+_{\varphi_i} - P_{\overline{\mathcal{O}}}$ for $i = 1, 2$ (see §6). The X-coordinate of the section $P^{(2)}_{\varphi_1}$ is given by

$$X(P^{(2)}_{\varphi_1}) = \frac{-(1+\sqrt{-1})}{(2(1+2\sqrt{-1})^2)}\Big(T^2 + c_1T + c_0\Big),$$

where

$$T = t + \sqrt{-1}/t,$$
$$c_1 = \sqrt{1-\sqrt{2}}\Big(9 + 13\sqrt{-1} - 2\sqrt{2} + 11\sqrt{-2}\Big), \quad c_0 = \frac{1}{6}\big(161 - 97\sqrt{-1}\big).$$

Since $P_{\overline{\mathcal{O}}}$ is defined over $k = \mathbf{Q}(\sqrt{2})$, $P^{(2)}_{\varphi_2}$ is also the image of $P^{(2)}_{\varphi_1}$ by the complex conjugate $\sqrt{-1} \mapsto -\sqrt{-1}$.

The explicit formula for the sections R_{22} described in Proposition 6.1 is

$$R_{22} = \Big((1-2\sqrt{-1})(2+\sqrt{2}-\sqrt{-2})\sqrt{1-\sqrt{2}} - \frac{(1-2\sqrt{-1})^4}{6}, T\Big).$$

Let $\sigma \in \mathrm{Gal}(L/\mathbf{Q})$ be the automorphism given by $\sqrt{1-\sqrt{2}} \mapsto -\sqrt{1-\sqrt{2}}$, and γ the complex conjugate $\sqrt{-1} \mapsto \sqrt{-1}$. Then, we have

$$R_{33} = -\sigma(R_{22}), \quad R_{23} = \gamma(R_{22}), \quad R_{32} = \gamma(R_{33}).$$

The height matrix with respect to $P^{(2)}_{\varphi_1}$, $P^{(2)}_{\varphi_2}$, R_{22}, R_{33}, R_{23} and R_{32} is given by

$$\frac{1}{3}\begin{pmatrix} 12 & 3 & 0 & 0 & -3 & -3 \\ 3 & 12 & -3 & -3 & 0 & 0 \\ 0 & -3 & 4 & 2 & 0 & 0 \\ 0 & -3 & 2 & 4 & 0 & 0 \\ -3 & 0 & 0 & 0 & 4 & 2 \\ -3 & 0 & 0 & 0 & 2 & 4 \end{pmatrix}.$$

Its determinant is $2^7/3^2$ as expected.

7.3. **The singular $K3$ surface $X_{[3,0,3]}$.** The transcendental lattice of the singular $K3$ surface $X_{[3,0,3]}$ is given by the matrix

$$\begin{pmatrix} 6 & 0 \\ 0 & 6 \end{pmatrix}.$$

In this case, complex elliptic curves $E_1/\mathbf{C}$ and $E_2/\mathbf{C}$ are given by

$$E_1 : \mathbf{C}/\mathbf{Z} \oplus \mathbf{Z}(3\sqrt{-1}), \quad E_2 : \mathbf{C}/\mathbf{Z} \oplus \mathbf{Z}\sqrt{-1}.$$

The j-invariants of them are given by

$$j(E_1) = 76771008 + 44330496\sqrt{3}, \quad j(E_2) = 1728.$$

We work with the base field $k = \mathbf{Q}(\sqrt{3})$, and we take the following Weierstrass forms, which corresponds to the formula (5.1) with $a = 9(2+\sqrt{3})$, $b = (1-\sqrt{3})/3$. (The equation of E_2 is scaled differently from (5.1).)

$$\begin{aligned} E_1 : \quad y_1^2 &= x_1^3 - (2+\sqrt{3})(3x_1 + 1 - \sqrt{3})^2 \\ E_2 : 27y_2^2 &= x_2^3 - 9(2+\sqrt{3})(x_2 - 3 - \sqrt{3})^2 \end{aligned}$$

E_1 and E_2 are isogenous with complex multiplication by $\mathbf{Q}(\sqrt{-1})$, and there are two 3-isogenies from E_1 to E_2. One is obtained by a coordinate changes of the 3-isogeny φ in §4. We denote it by φ_1. The other, denoted by φ_2, is a composite of φ_1 and the complex multiplication $[\sqrt{-1}]$ on E_2. With our equation of E_2, the multiplication-by-(-1) map on E_2 is given by $(x_2, y_2) \mapsto (-x_2 + 12 + 6\sqrt{3}, \sqrt{-1}y_2)$.

The 3-isogenies φ_1, φ_2 are given by

$$\begin{aligned} \varphi_1(x_1, y_1) &= (\varphi_x(x_1), \varphi_y(x_1)y_1)\,, \\ \varphi_2(x_1, y_1) &= \left(-\varphi_x(x_1) + 12 + 6\sqrt{3}, \sqrt{-1}\varphi_y(x_1)y_1\right), \end{aligned}$$

where

$$\begin{aligned} \varphi_x(x_1) &= \frac{x_1^3 + 12(2+\sqrt{3})x_1^2 + 12(1+\sqrt{3})x_1 + 8}{3x_1^2}, \\ \varphi_y(x_1) &= \frac{\left(x_1 + 2(1+\sqrt{3})\right)\left(x_1^2 - 2(1+\sqrt{3})x_1 + 4(1-\sqrt{3})\right)}{27x_1^3}. \end{aligned}$$

Over the base field k, the equation of $F^{(6)}_{E_1,E_2}$ is given by

$$F^{(6)}_{E_1,E_2} : Y^2 = X^3 - (387 + 224\sqrt{3})X + 3(3 + 2\sqrt{3})u^6 + \frac{45 + 26\sqrt{3}}{9u^6}.$$

By letting $s' = s(3+2\sqrt{3})/9 = u^6(3+2\sqrt{3})/9$, the equation of $F^{(1)}_{E_1,E_2}$ is given by

$$F^{(1)}_{E_1,E_2} : Y^2 = X^3 - (387 + 224\sqrt{3})X + (7 + 4\sqrt{3})\Big(s' + \frac{1}{s'}\Big).$$

Using the method described in §5, the section $P^{(1)}_{\varphi_1}$ of $F^{(1)}_{E_1,E_2}$ is given by

$$P^{(1)}_{\varphi_1} = \left(\frac{c_3S'^3 + c_2S'^2 + c_1S' + c_0}{144(S'+2)}, \frac{s'(s'-1)\big(d_4S'^4 + d_3S'^3 + d_2S'^2 + d_1S' + d_0\big)}{1728\,(s'+1)^3}\right),$$

where

$$
\begin{aligned}
&S' = s' + \frac{1}{s'},\\
&c_3 = (2-\sqrt{3})^2, \quad c_2 = -42, \quad c_1 = 12(91+36\sqrt{3}), \quad c_0 = -8(1267+680\sqrt{3}),\\
&d_4 = (2-\sqrt{3})^3, \quad d_3 = -4(25-12\sqrt{3}), \quad d_2 = 24(107+15\sqrt{3}),\\
&d_1 = 16(461+444\sqrt{3}), \quad d_0 = -16(54676+32091\sqrt{3}).
\end{aligned}
$$

If we write $P^{(1)}_{\varphi_1} = (x(s'), y(s'))$, then we have $P^{(1)}_{\varphi_2} = (-x(-s'), \sqrt{-1}\, y(-s'))$. The sections $P^{(1)}_{\varphi_1}$ and $P^{(1)}_{\varphi_2}$ form a basis of $F^{(1)}_{E_1,E_2}(\bar{k}(s)) = F^{(1)}_{E_1,E_2}(\mathbf{Q}(\sqrt{3},\sqrt{-1})(s))$. The height matrix with respect to this basis is given by

$$
\begin{pmatrix} 6 & 0 \\ 0 & 6 \end{pmatrix}.
$$

Next, we consider $F^{(2)}_{E_1,E_2}(\bar{k}(t))$. The field of definition of 2-torsion subgroups $E_1[2]$ and $E_2[2]$ is $\mathbf{Q}(\sqrt{3+2\sqrt{3}})$. In order to compute generators of $F^{(2)}_{E_1,E_2}(\bar{k}(t))$, we need to work with the field $L = \mathbf{Q}\big(\sqrt{3+2\sqrt{3}}, \sqrt{-1}\big)$.

The explicit formulas for the sections $P^{(2)}_{\varphi_1}$, $P^{(2)}_{\varphi_2}$, R_{22}, R_{33}, R_{23}, R_{32} described in Proposition 6.1 are as follows:

$$
\begin{aligned}
P^{(2)}_{\varphi_1} &= \Big(\frac{1}{6}\Big(\sqrt{3}\, T_+^2 + 6(1+\sqrt{3})\, T_+ + 42 + 20\sqrt{3} \Big),\\
&\qquad\qquad \frac{\alpha}{12} T_- \Big((1-\sqrt{3})\, T_+^2 - 6\sqrt{3}\alpha\, T_+ + 112 + 56\sqrt{3} \Big) \Big)\\
P^{(2)}_{\varphi_2} &= \Big(\frac{1}{6}\Big(\sqrt{3}\, T_-^2 + 6\sqrt{-1}(1+\sqrt{3})\, T_- - 42 - 20\sqrt{3} \Big),\\
&\qquad\qquad -\frac{\alpha}{12} T_+ \Big(-(1-\sqrt{3})\, T_+^2 - 6\sqrt{-3}\alpha\, T_+ - 112 - 56\sqrt{3} \Big) \Big)\\
R_{22} &= \Big(-7 - 4\sqrt{3} + 2(1+\sqrt{3})\alpha, (2+\sqrt{-3}) T_+ \Big),\\
R_{33} &= \Big(-7 - 4\sqrt{3} - 2(1+\sqrt{3})\alpha, -(2+\sqrt{-3}) T_+ \Big),\\
R_{23} &= \Big(7 + 4\sqrt{3} - 2(1+\sqrt{3})\alpha, (2+\sqrt{-3}) T_- \Big),\\
R_{32} &= \Big(7 + 4\sqrt{3} + 2(1+\sqrt{3})\alpha, -(2+\sqrt{-3}) T_- \Big),
\end{aligned}
$$

where

$$
\alpha = \sqrt{3+2\sqrt{3}}, \quad T_+ = \frac{3t}{\alpha} + \frac{\alpha}{3t}, \quad T_- = \frac{3t}{\alpha} - \frac{\alpha}{3t}.
$$

The height matrix with respect to $P^{(2)}_{\varphi_1}$, $P^{(2)}_{\varphi_2}$,R_{22}, R_{33}, R_{23}, R_{32} is given by

$$
\frac{1}{3}\begin{pmatrix}
12 & 0 & 0 & 0 & -3 & -3 \\
0 & 12 & -3 & -3 & 0 & 0 \\
0 & -3 & 4 & 2 & 0 & 0 \\
0 & -3 & 2 & 4 & 0 & 0 \\
-3 & 0 & 0 & 0 & 4 & 2 \\
-3 & 0 & 0 & 0 & 2 & 4
\end{pmatrix}.
$$

Its determinant is 2^4 as expected.

References

[1] Hiroshi Inose, *Defining equations of singular $K3$ surfaces and a notion of isogeny*, Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., Kyoto, 1977), Kinokuniya Book Store, Tokyo, 1978, pp. 495–502. MR 578868

[2] A. Kumar and M. Kuwata, *Elliptic K3 surfaces associated with the product of two elliptic curves: Mordell-Weil lattices and their fields of definition*, arXiv:1409.2931, to appear in Nagoya Math. J.

[3] Masato Kuwata, *The canonical height and elliptic surfaces*, J. Number Theory **36** (1990), no. 2, 201–211. MR 1072465

[4] ———, *Elliptic $K3$ surfaces with given Mordell-Weil rank*, Comment. Math. Univ. St. Paul. **49** (2000), no. 1, 91–100. MR 1777156

[5] Masato Kuwata and Tetsuji Shioda, *Elliptic parameters and defining equations for elliptic fibrations on a Kummer surface*, Algebraic geometry in East Asia—Hanoi 2005, Adv. Stud. Pure Math., vol. 50, Math. Soc. Japan, Tokyo, 2008, pp. 177–215. MR 2409557

[6] T. Shioda and H. Inose, *On singular $K3$ surfaces*, Complex analysis and algebraic geometry, Iwanami Shoten, Tokyo, 1977, pp. 119–136. MR 0441982

[7] Tetsuji Shioda, *Kummer sandwich theorem of certain elliptic $K3$ surfaces*, Proc. Japan Acad. Ser. A Math. Sci. **82** (2006), no. 8, 137–140. MR 2279280

[8] ———, *Correspondence of elliptic curves and Mordell-Weil lattices of certain elliptic $K3$'s*, Algebraic cycles and motives. Vol. 2, London Math. Soc. Lecture Note Ser., vol. 344, Cambridge Univ. Press, Cambridge, 2007, pp. 319–339. MR 2385296

[9] ———, *The Mordell-Weil lattice of $y^2 = x^3 + t^5 - 1/t^5 - 11$*, Comment. Math. Univ. St. Pauli **56** (2007), no. 1, 45–70. MR 2356749

[10] Jaap Top, *Descent by 3-isogeny and 3-rank of quadratic fields*, Advances in number theory (Kingston, ON, 1991), Oxford Sci. Publ., Oxford Univ. Press, New York, 1993, pp. 303–317. MR 1368429

Faculty of Economics, Chuo University
742-1 Hachioji-shi, Tokyo 192-0393 Japan
E-mail address: kuwata@tamacc.chuo-u.ac.jp

College of Science and Engineering, Ritsumeikan University
1-1-1 Noji-higashi, Kusatsu Shiga 525-8577 Japan
E-mail address: kutsumi@fc.ritsumei.ac.jp